



Utilization of complementary and alternative medicine for the prevention of COVID-19 infection in Ghana: A national cross-sectional online survey

Irene A. Kretchy^a, Joseph A. Boadu^b, James-Paul Kretchy^c, Kofi Agyabeng^d, Alfred A. Passah^b, Augustina Koduah^a, Kwabena F.M. Opuni^{b,*}

^a Department of Pharmacy Practice and Clinical Pharmacy, School of Pharmacy, University of Ghana, P.O. Box LG43, Legon, Accra, Ghana

^b Department of Pharmaceutical Chemistry, School of Pharmacy, University of Ghana, P.O. Box LG43, Legon, Accra, Ghana

^c Department of Physician Assistantship Studies, School of Medicine and Health Sciences, Central University, Miotso, Accra, Ghana

^d Department of Mathematics, KU Leuven, Leuven, Belgium

ARTICLE INFO

Keywords:

CAM use
Coronavirus
COVID-19
Herbs
Perception
Side effects

ABSTRACT

Complementary and alternative medicine (CAM) use is widespread and has played critical roles in preventing infections, including previous coronaviruses. This study sought to document current practices in the use of CAM for the prevention of COVID-19 disease in Ghana. An anonymous electronic survey was conducted from February 1, 2021 to April 30, 2021. Data on demographic characteristics, basic clinical information, illness perceptions about COVID-19, and CAM use during the pandemic period were generated. While about 82.5% (986/1195) of the participants used CAM during the COVID-19 period, 69.1% (681/986) of CAM users intended it for COVID-19 infection prevention. Vitamin supplements (88.1%, 869/986), spiritual healing/prayer (23.3%, 230/986), mineral supplements (22.3%, 220/986), botanical/herbal medicines (22.2%, 219/986), and diet therapy (19.4%, 191/986) were the main types of CAM used. From the adjusted binary logistic regression model, current age (aOR: 1.03, 95%CI: 1.01–1.05), sex (aOR: 1.41, 95%CI: 1.02–1.95), participants' perceptions of consequences (aOR: 1.10, 95%CI: 1.04–1.17), identity (aOR: 1.15, 95%CI: 1.06–1.25) and concerns about COVID-19 (aOR: 0.91, 95%CI: 0.85–0.97) were statistically significant predictors of CAM use. These results suggest the need for appropriate public health policy on COVID-19 and CAM use in addition to directing further research initiatives toward an optimized COVID-19 prevention scheme using clinically validated CAM treatments. Research to validate the clinical efficacy of these products, especially the herbs, for COVID-19 prevention while isolating lead compounds that could be optimized and used for the treatment and prevention of COVID-19 is also recommended.

1. Introduction

The Corona Virus Disease (COVID-19) which is a zoonotic viral infectious disease caused by severe acute respiratory syndrome coronavirus 2 has become a global pandemic of major public health concern (Gorbalenya et al., 2020). Worldwide, over 200 million cases and more than 4 million COVID-19 related deaths have been reported (COVID-19 Dashboard by the Center for Systems Science and Engineering (CSSE), 2021) with individuals having compromised immune systems, chronic diseases, and the elderly as high-risk populations (Wu and McGoogan, 2020). Although the World Health Organization developed preventive guidelines to slow viral spread (Wu and McGoogan, 2020), Medicine Regulatory Agencies in several countries have approved vaccines under

emergency use authorisation and medicines like azithromycin, hydroxychloroquine, chloroquine phosphate, ivermectin, doxycycline, dexamethasone, methylprednisolone, remdesivir, and convalescent plasma for COVID-19 case management (Tarighi et al., 2021; Sanders et al., 2020).

With these conventional interventions, there are reports of complementary or alternative treatments such as herbal preparations, dietary therapy, vitamin supplements, and prayer for prevention or treatment of COVID-19 (Jabaris and Ananthlakshmi, 2021; Boozari and Hosseinzadeh, 2020; Panyod et al., 2020).

CAM involves medications, treatments, and medicinal practices used concurrently with (complementary) or in place of (alternative) conventional treatments (Ernst and Fugh-Berman, 2002; Lichtenstein and

* Corresponding author.

E-mail address: kfopuni@ug.edu.gh (K.F.M. Opuni).

<https://doi.org/10.1016/j.pmedr.2021.101633>

Received 21 August 2021; Received in revised form 28 September 2021; Accepted 7 November 2021

Available online 9 November 2021

2211-3355/© 2021 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Waaen, 2002). Utilization of CAM in several countries is high (Harris et al., 2012; Kretchy et al., 2014). A review reporting CAM use in Sub-Saharan Africa ranged from 4.6% to 94% with an average of 58.2% (James et al., 2018). High availability, easy access, low costs of CAM (Kretchy et al., 2014; James et al., 2018; Debas et al., 2006), and perception towards illness (Canaway and Manderson, 2013; Kucuk, 2017) mainly contribute to its expanded utilization.

How people perceive illness is key to how they adopt particular health behaviours for prevention or management (Figueiras and Neto, 2019). Illness perceptions are usually cognitive or emotional representations of a disease that are predictive of coping (Broadbent et al., 2006) and based on the common-sense self-regulatory theory of illness representations (Leventhal et al., 1980). This theory proposes that people create mental representations of illness experiences through cognitive, and affective processes based on available information or symptom experiences (Broadbent et al., 2006). The theory proposes five components of cognitive illness perceptions, namely consequences, timelines, personal control, treatment control, and identity, while the emotional illness perception is based on concerns and negative emotional arousal of the illness (Broadbent et al., 2006; Leventhal et al., 2016).

Illness perception has successfully predicted behaviours and outcomes across populations including hypertension, diabetes, coronary heart disease, anxiety, depression, cancer and medication adherence (Foxwell et al., 2013; Broadbent et al., 2015; Alyami et al., 2021; Anakwa et al., 2021). In recent times, studies have explored illness perception and adherence to safety protocols for COVID-19 and observed that perceptions toward COVID-19 had significant impact on adherence to these measures (Chong et al., 2020). The physical and psychological impact of illness perception during the COVID-19 pandemic as well as the role of CAM and integrative therapies as coping health behaviours in response to perceptions about COVID-19 have also been reported (Kristoffersen et al.; Man et al., 2020; Chong et al., 2021; Skapinakis et al., 2020; Dias Neto et al., 2021).

Natural medicines have played essential roles in previous coronavirus infection prevention (Boozari and Hosseinzadeh, 2020) and threats of the current COVID-19 with its associated risk perceptions has led to CAM utilization as well (Paudyal et al., 2021). Herbs such as *Allium sativum*; *Camellia sinensis*; *Zingiber officinale*, *Nigella sativa*, *Glycyrrhiza glabra*, and *Astragalus membranaceus* were used for preventing previous coronavirus infections (Luo et al., 2020) with vitamin and mineral supplements reported for COVID-19 prevention (Gasmi et al., 2021; Kretchy et al., 2021).

CAM use in the treatment and management of diseases is prevalent (Kretchy et al., 2014; Kretchy et al., 2021; Kuunibe and Domanban, 2012; Yarney et al., 2013), yet information on COVID-19 perception influencing CAM is limited. This study therefore aimed to assess the role of illness perceptions towards COVID-19 and CAM utilization while documenting the prevalence and pattern of use. Understanding these relationships will inform further research initiatives toward optimized COVID-19 prevention using clinically validated CAM treatments.

2. Methods

The study was approved by the Institutional Review Board of the Noguchi Memorial Institute for Medical Research, University of Ghana (CPN: 028/20-21) and conducted in accordance with the Helsinki Declaration. Participants were voluntarily recruited after they had been informed about the purpose of the study. All study participants approved online written informed consent before commencement of data collection.

2.1. Study design and participants

This was a national cross-sectional online study using anonymous electronic survey from February to April 2021 via an online platform (<https://ee.kobotoolbox.org/single/2q6zUPIK>). The survey tool was

circulated through emails, professional association groups, and five social media platforms – Telegram, WhatsApp, Instagram, LinkedIn, and Facebook. Due to the threatening second wave of COVID-19 and risks associated with conducting face-to-face community-based national surveys, the data collection process was online, and participants were recruited across all sixteen regions of Ghana using a mix of convenience and snowball sampling approaches to increase the number of participants.

Participants were Ghanaian residents aged 18 years or more, able to understand the questions in English, and had access to internet. Screening questions to ascertain age and residency status of participants were used.

Participation was voluntary after brief information on study objectives, confidentiality, and estimated completion time were provided. Participants had access to the link after informed consent was obtained. Participants were requested to share the invitation with their contacts. To control for multiple submissions, the survey settings were set to reject multiple responses from the same IP address.

2.2. Data collection

A 3-paged 30-item questionnaire was used to generate data on demographic characteristics, basic clinical information, illness perceptions about COVID-19, and CAM use during the pandemic (See Fig. 2 for complete list of CAM therapies). Demographic and clinical questions included sex, age, region of residence, educational status, existing health condition and COVID-19 risk vulnerability. Questions on COVID-19 tests and outcomes were indicated. The frequency, type, patterns, sources, reasons, beliefs, and side effects of CAM use was also noted. Examples of questions asked included (1) *Which of the following CAM have you used during the COVID-19 pandemic period?* (2) *Was the CAM intended to prevent COVID-19 infection?* (3) *How long have you been using the CAM during the COVID-19 pandemic?* and (4) *Where did you receive the information on CAM?* Participants' perceptions about COVID-19 were assessed with the 9-item Brief Illness Perception Questionnaire (Broadbent et al., 2006). The items are scored on a 10-point Likert scale (except for item 9) for consequences, timeline, personal control, treatment control, identity, concern, illness understanding, and emotional response dimensions to COVID-19 perceptions. Item 9, an open-ended question on beliefs about causes of illness was excluded from this study, and in line with previous research, the overall score for the B-IPQ and Cronbach's alpha were not computed because each subscale is measured by only one item (Anakwa et al., 2021; Haines et al., 2019).

To minimize non-response and high drop-out rates, the questionnaire was kept as short as possible, and participants could answer all questions within 5 min. The adaptive questioning approach was also applied to minimize the response time of participants by conditionally showing specific questions based on responses to previous items.

An online pre-test of the questionnaire was conducted among 20 people who were not members of the research team and played no role in the design of the questionnaire, to ensure that the electronic version was user-friendly, technically functional, and comprehensible by prospective participants. Prior to piloting the online version, face-to-face interviews with consideration for strict adherence to the COVID-19 prevention protocols were conducted among 7 people to ensure that the questions were comprehensible and coherent.

2.3. Data analysis

Data collected through the electronic questionnaire were extracted into excel for cross-validation and cleaning and exported to STATA version 15 (StataCorp LLC, USA) for analysis. The cross-validation involved checking consistencies in responses and ensuring that all respondents met the inclusion criteria through the use of frequency and cross-tabulation of responses. Descriptive statistics of responses for categorical variables were reported with frequencies and percentages

Table 1
Background characteristics of study participants.

	Frequency	Percent
Current Age		
Median (LQ, UQ)	25 (21, 33)	
18–30	837	70.0
31–40	223	18.7
41–50	104	8.7
>=51	31	2.6
Sex		
Female	644	53.9
Male	551	46.1
Religion		
Christianity	1,117	93.5
Islam	69	5.8
Traditionalism	3	0.3
No religion	6	0.5
Educational Level		
Primary	6	0.5
Secondary	60	5.0
Tertiary	1,129	94.5
Tested for COVID-19		
No	863	72.2
Yes	332	27.8
Test results		
Negative	274	82.5
Positive	58	17.5

while that of continuous variables were reported in terms of means and standard deviations when normally distributed but reported as median with interquartile ranges when skewed. Bar charts were used to graphically display the distribution of existing health conditions of study participants, CAM used during the COVID-19 pandemic period, sources of information on CAM and sources of CAM among the study participants. Pearson Moment Correlation Coefficient was used in assessing the relationship between the various sub domains of perceptions about COVID-19 illness. A binary logistic regression model was used to assess the effects of background characteristics and illness perceptions about COVID-19 on the use of CAM. The results of the model were reported as odds ratios with their 95% confidence intervals. All statistical tests of significance were conducted at a 5% level.

3. Results

3.1. Background characteristics of study participants

This study had 1,195 participants in total. Participants had a median age of 25 years with more than two-thirds of them between ages 18 and 30 years inclusive. Few (2.6%, 31/1195) participants were 51 years and beyond. The majority (53.9%, 644/1195) were females. Almost all (93.5%, 1117/1195) participants were Christians and had tertiary level education (94.5%, 1129/1195). Less than a third (332/1195) of the participants had tested for COVID-19 out of which 17.5% (58/332) were positive (Table 1). Three common existing health conditions among participants were common cold (4.8%, 57/1195), asthma (2.9%, 35/1195), and hypertension (2.9%, 34/1195) (Fig. 1).

3.2. Pattern of CAM use among study participants

In all 82.5% (986/1195) of the participants used some form of CAM during the COVID-19 period. Five commonly used CAM were vitamin supplements (88.1%, 869/986), spiritual healing/prayer (23.3%, 230/986), mineral supplements (22.3%, 220/986), botanical/herbal medicines (22.2%, 219/986), and diet therapy (19.4%, 191/986). The nature of CAM was mainly raw material/homemade (21.8%, 215/986) and finished products (75.3%, 742/986) (Table 2). The raw materials/homemade preparations were reported for use either individually or as combination remedies (see Appendix A). The least used CAM were magnets therapy, chelation therapy, and therapeutic touch/reiki (Fig. 2). About 86.5% (853/986) reported no side effects to CAM (Table 2). Participants' source of information on CAM was mostly from health practitioners (46.9%, 462/1195) or Family/Friends/Colleagues (44.3%, 436/1195) (Fig. 3).

Medicines retail outlets (pharmacy and chemical shops) were the major sources of CAM acquisition while Herbal stores were the least (Fig. 4). The CAMs used were mostly finished products (75.3%, 742/986) and were registered by the Food and Drugs Authority of Ghana (73.2%, 722/986). About a third of participants had used CAM for more than six months (31.3%, 309/986) and 69.1% (681/986) of CAM use during this period being intended for COVID-19 infection prevention. More than half (58.2%, 574/986) of the participants took CAM daily (Table 2). The frequency of use was mostly based on personal decisions (48.5%, 478/986) with a few from social media. Similar proportions of




Fig. 1. Existing health conditions of study participants.

Table 2
Pattern of CAM use among study participants.

	Frequency	Percent
CAM use		
No	209	17.5
Yes	986	82.5
Nature of the CAM		
Raw materials/home-made	215	21.8
Finished product	742	75.3
Other (specify)	29	2.9
CAM registered by the FDA		
Yes	722	73.2
No	76	7.7
Don't Know	105	10.7
Not Applicable	83	8.4
Duration for using the CAM during the COVID-19		
<1 month	252	25.6
1–3 months	303	30.7
4–6 months	122	12.4
More than 6 months	309	31.3
How did you decide on the frequency of therapy		
Followed product label instructions	267	27.1
Followed practitioner's instructions	224	22.7
Decided myself	478	48.5
Media (e.g. TV, Radio, Newspapers)	99	10.0
Friends/Relatives	157	15.9
Social media	63	6.4
Internet	83	8.4
Other	10	1.0
CAM intended to prevent COVID-19 infection		
Yes	681	69.1
No	305	30.9
How often did you use the CAMs		
Daily	574	58.2
Weekly	242	24.6
Monthly	88	8.9
Annually	13	1.3
Biannually	2	0.2
Other	67	6.8
CAMs are effective in preventing COVID-19 infections		
Strongly Agree	182	18.5
Agree	439	44.5
Neutral	304	30.8
Disagree	44	4.5
Strongly Disagree	17	1.7
Safe to use CAM in preventing COVID-19 infection		
Strongly Agree	148	15.0
Agree	478	48.5
Neutral	296	30.0
Disagree	55	5.6
Strongly Disagree	9	0.9
Side effects experienced^{a)}		
Headache	33	3.35
Dizziness	25	2.54
Diarrhoea	23	2.33
Fatigue	21	2.13
Nausea	20	2.03
Allergic reactions	17	1.72
Vomiting	10	1.01
Pain	7	0.71
Bleeding	1	0.1
Bruising skin	1	0.1
Dermatitis	1	0.1

^{a)}Multiple responses.

CAM users agreed or strongly agreed to CAM being effective (63%, 621/986) and safe (63.5%, 626/986) for use in COVID-19 infection prevention (Table 2).

3.3. Perceptions about COVID-19 among study participants

COVID-19 illness representations included consequences of the infection (4.61 + 2.96), timeliness of living with the illness (5.23 + 2.93), personal control over the virus (4.84 + 2.78), identity with experience of COVID-19 symptoms (1.65 + 2.30) and concerns about

COVID-19 (7.45 + 2.80) (Table 3).

3.4. Effects of background characteristics and perceptions about COVID-19 on the use of CAM

From the binary logistic regression model, current age, sex, participants' perceptions on consequence, identity, and concern about COVID-19 were the only statistically significant predictors of CAM use (Table 4).

The odds of using CAM increased by 3% for each year increase in age (aOR: 1.03, 95%CI: 1.01–1.05). Female participants had 41% higher odds of using CAM compared with their male counterparts (aOR: 1.41, 95%CI: 1.02–1.95).

Each unit increase in participants' perception of the consequences of COVID-19 led to a 10% increase in the odds of using CAM (aOR: 1.10, 95%CI: 1.04–1.17). A unit increase in beliefs about COVID-19 symptom experience was associated with a 15% increase in the odds of using CAM (aOR: 1.15, 95%CI: 1.06–1.25). The odds of using CAM reduced by 9% with every unit decrease in participants' level of concern about COVID-19 (aOR: 0.91, 95%CI: 0.85–0.97).

4. Discussion

This study investigated uptake of CAM for prevention of COVID-19 infection in Ghana to understand the trend and suggest context-specific alternatives to the management of the disease. A preponderance of our study participants were females, a finding that corresponded to national statistics where females formed a majority of 51.2% with higher life expectancy of 65 years, compared to 63 years in males (2010 Population Housing Census: National Analytical Report, 2021). The majority of our study participants had attained tertiary education and this is common with on-line studies where people who are literates can respond to such surveys (Hossain et al., 2020; Nekliudov et al., 2020). The study also recorded highest number of participants as Christians and this corroborates the report that about 70% of the Ghanaian population is estimated to be Christians (2010 Population Housing Census: National Analytical Report, 2021). Ghana and other Sub-Saharan African countries, the evidence of using faith-based approaches for health problems in combination with CAM has been reported (Gyasi et al., 2015; Okoronkwo et al., 2014).

In this study, less than a third of participants had tested for COVID-19 of which 17.5% were positive. This low rate of testing for COVID-19 confirms the low national trend in which approximately seven tests per 100, 000 people were recorded daily (Ghana's Outbreak Response Management Updates, 2021). This portrays deficiencies in existing testing facilities and the lack of capacity to accommodate testing needs of the population of about 30 million, should there be a sudden increase in COVID-19 cases in the country. It is hoped that the government would improve testing capacity, by adequately equipping laboratories, since early detection of cases impact disease management and prevents spread.

Our findings also showed that common cold, asthma, and hypertension were common health conditions reported. It is known that whilst common cold/flu and fever may be experienced in the early stages of COVID-19, among other respiratory symptoms like cough and dyspnea (Ahmad et al., 2021), asthma, diabetes, and hypertension may complicate the outcome of coronavirus infections (Yawson et al., 2020).

The products used by the respondents were consistent with most frequently used CAM products in literature (Kretchy et al., 2021; Okoronkwo et al., 2014; Gyasi et al., 2015; Egede et al., 2002). While about 69.1% of CAM use was intended for the prevention of COVID-19 disease, they were generally considered effective and safe for the anticipated purpose. Vitamins were the most frequently used to prevent COVID-19 followed by spiritual healing/prayers, use of mineral supplements, and botanical/herbal medicines. High use of CAM products has been linked with medical conditions considered as life-threatening (Egede et al., 2002). Since CAM was reported to be useful in




Fig. 2. CAMs used during the COVID-19 pandemic period by study participants.




Fig. 3. Source of Information on CAM.



Fig. 4. Sources of CAM among the study participants.

improving immunity in individuals, it has been suggested for use as COVID-19 prophylaxis. Traditional Chinese Medicine and Ayurveda have been used for COVID-19 infection prevention and/or treatment in countries like China and India (National Health Commission of the

People's Republic of China, 2021; Protocol, 2021). While dietary therapy and herbal medicines have also been suggested as complementary preventive therapy for COVID-19 (Panyod et al., 2020), a survey conducted in Iran revealed that the participants used herbal products, traditional medicines, and vitamin supplements to prevent them from

Table 3
Perceptions about COVID-19 among study participants.

	Mean + SD	Pearson Moment Correlation Coefficient							
		Consequence	Timeline	Personal control	Treatment control	Identity	Concern	Understanding	Emotional response
Consequence	4.61 + 2.96	1							
Timeline	5.23 + 2.93	0.34*	1						
Personal control	4.84 + 2.78	0.04	-0.04	1					
Treatment control	4.41 + 2.35	-0.06*	-0.08*	0.28*	1				
Identity	1.65 + 2.30	0.16*	0.13*	-0.05	-0.07*	1			
Concern	7.45 + 2.80	0.21*	0.18*	0.01	-0.13*	-0.01	1		
Understanding	2.25 + 2.42	-0.06*	-0.12*	0.18*	0.17*	0.05	-0.27*	1	
Emotional response	4.62 + 2.98	0.34*	0.17*	0.08*	-0.05	0.18*	0.33*	-0.03	1

SD: Standard Deviation, *P-value < 0.05.

Table 4
Effects of background characteristics and perceptions about COVID-19 on the use of CAM.

	Unadjusted Binary Logistic Regression Model			Adjusted Binary Logistic Regression Model		
	uOR	95%CI	P-value	aOR	95%CI	P-value
Current Age	1.04	1.02–1.06	<0.001***	1.03	1.01–1.05	0.005**
Sex			0.190			0.036*
Male	1.00			1.00		
Female	1.22	0.91–1.65		1.41	1.02–1.95	
Religion			0.680			0.602
Christian	1.13	0.63–2.03		1.18	0.64–2.18	
other	1.00			1.00		
Educational Level			0.250			0.706
Non-tertiary	1.00			1.00		
Tertiary	1.42	0.78–2.57		1.13	0.6–2.15	
Existing Health Conditions			<0.001***			0.064
No	1.00			1.00		
Yes	0.46	0.28–0.75		0.62	0.37–1.03	
Perceptions about COVID-19						
Consequence	1.14	1.08–1.2	<0.001***	1.10	1.04–1.17	0.001**
Timeline	1.07	1.02–1.13	0.010*	1.02	0.96–1.07	0.580
Personal control	0.97	0.92–1.03	0.320	0.99	0.93–1.05	0.711
Treatment control	0.95	0.89–1.01	0.130	0.96	0.9–1.03	0.299
Identity	1.21	1.11–1.31	<0.001***	1.15	1.06–1.25	0.001**
Concern	0.97	0.92–1.03	0.300	0.91	0.85–0.97	0.003**
Understanding	0.99	0.93–1.05	0.660	0.98	0.92–1.05	0.563
Emotional response	1.09	1.04–1.15	<0.001***	1.05	0.99–1.12	0.079

*p-value < 0.05, **p-value < 0.01, ***p-value < 0.001, CI: Confidence interval, aOR: Adjusted odds ratio, uOR: unadjusted odds ratio.

contracting the virus (Erfani et al., 2020). Some of the herbs used as homemade CAM intervention in this study are in line with previous reports (Paudyal et al., 2021; Luo et al., 2020). The use of spiritual therapy is similar to other studies, which reported that prayer intervention was one of the common forms of CAM used among residents in Ghana and South African (Gyasi et al., 2015; Singh et al., 2004). Prayers are highly favoured within most African communities and may account for the high numbers of respondents resorting to this form of CAM (Okoronkwo et al., 2014; Gyasi et al., 2015; Singh et al., 2004). Most respondents sourced their CAM from medicines retail outlets (pharmacies and over-the-counter medicine sellers), and this is supported by previous work, in which the pharmacy was an important source for CAM (Gyasi et al., 2015). The medicine retail outlets are usually the first point of call for medical care (Okai et al., 2019). The majority of the respondents used CAM products because of personal decisions. Having a strong personal belief has been documented to play a critical role in individuals' use of CAM (Gyasi et al., 2016). Based on the participants' reports, and in line with a previous study on the use of herbal medicinal products (Kretchy et al., 2021), the majority of the CAM products were registered by the national Food and Drugs Authority (FDA), confirming a rise in the number of registered products by FDA Ghana from 2011 to 2019 (Kretchy et al., 2021) though not specific for COVID-19.

The demographic characteristics that significantly influenced the trend of CAM use were the age and sex of the participants. The findings

indicated that the use of CAM during the COVID-19 increased with the increasing age of participants. Although previous studies on age and CAM use have reported mixed results, a review found that CAM use significantly increased with increasing age in 26 studies out of the 134 that were analysed (Bishop and Lewith, 2010). In relation to the findings on sex, this study observed that female participants had a higher chance (41%) of using CAM than males during the COVID-19 pandemic. Similar results have been documented for females being more likely to use CAM than males (Egede et al., 2002; Alwhaibi et al., 2015; Laiyemo et al., 2015; Alwhaibi and Sambamoorthi, 2016), which are congruent with our findings. Plausible reasons for this observation are that women are more motivated to use CAM because they tend to adopt more preventive healthcare approaches, as well as have more health needs that may not be sufficiently satisfied by conventional healthcare systems (Zhang et al., 2015; Kristoffersen et al., 2014).

The perceptions of people about COVID-19 will be relevant to how they prevent or adjust to the illness. These cognitive and emotional representations of COVID-19 have been reported to be formed at the initial stages of the pandemic and have remained quite stable or decreased over time (Dias Neto et al., 2021). In this study, participants formed their perceptions about COVID-19 based on both cognitive and emotional views about the disease which comprises the anticipated physical, emotional and social effects and outcomes of the illness on an individual's wellbeing (Broadbent et al., 2006). Our study revealed that

the participants' perception of COVID-19 affecting their lives (consequences) led to the use of CAM. COVID-19 has resulted in the significant loss of many lives globally, an increase in unemployment rates, a reduction in food security as well as self-isolation of infected individuals (Piltch-Loeb et al., 2021; Aliakbari Dehkordi et al., 2020; VanderWeele, 2020; Yang et al., 2020; Mentis, 2021). These consequences are likely reasons for people's use to CAM for COVID-19 prevention. Our study indicated that the beliefs about experiencing COVID-19 symptoms also resulted in a significant increase in CAM use. Some common symptoms of COVID-19 include fever, cough and dyspnoea (Pullen et al., 2020). However, there are more severe symptoms of COVID-19 such as breathlessness (from respiratory distress syndrome), sudden confusion, and constant chest pain (Valizadeh et al., 2020; Ahmad et al., 2020). The participants may have experienced any of these symptoms or want to avoid experiencing such symptoms and thus, resulting in their use of CAM in preventing COVID-19 infections. The results of this study also showed that the lower the concern about COVID-19, the lower the likelihood of using CAM. In the course of a pandemic, fear, worry, or even paranoia can easily spread throughout the population (Freckelton, 2020). This fear tends to push individuals to take drastic measures in avoiding infection. These measures include using readily available and affordable health interventions, such as herbal medicine and dietary supplements (Panyod et al., 2020; Freckelton, 2020; Alyami et al., 2020; Silveira et al., 2020). Thus, the lower the concerns, the less likely to take any action.

The strength of this study is in the use of an online survey that enabled the recruitment of study participants across all the regions of Ghana since the COVID-19 pandemic is a challenge to conducting face-to-face research. Also, this study provides comprehensive information on the prevalence, pattern, and perceptions towards CAM use for COVID-19 pandemic control in Ghana, which can serve as the basis for an improved COVID-19 management scheme using clinically validated CAM treatments.

There are however some limitations that need to be considered in the interpretation of the study findings. First, online data collection was used for this study and therefore only people who could read and write, have access to social media networks and access the internet could participate. Second, since convenience and snowball sampling were used in this study, the data may not be representative of the population distribution in Ghana in terms of age, sex, and place of residence. Hence, a disproportionately large number of study participants were female and had tertiary education, which is a common observation in online surveys (Hossain et al., 2020; Nekliudov et al., 2020). Third, the cross-sectional nature of this study prevents the analysis of causal relationships, and the use of self-reported measures introduces reporting bias. Finally, in considering the prevalence, pattern, and perceptions towards the COVID-19 pandemic and CAM use for its control, the long-term effects of CAM cannot be inferred from this study.

Despite these limitations, this study findings highlight some key clinical, research, and policy implications. The outcome of the study indicates a high prevalence of CAM use for COVID-19 prevention. Consequently, there is a clinical implication of potential herb-drug interaction for people who are on medications for chronic disease conditions. Thus, a rigorous health educational campaign is needed to avoid any potential public health crisis. In addition, a clear policy is needed for the use of CAM in the prevention and management of COVID-19. From this study, many people used different herbs for the prevention of COVID-19 infection and as a result, research could be conducted to validate the clinical efficacy of these herbs for COVID-19 prevention and possibly isolate lead compounds that could be optimized and used for the treatment and prevention of the coronavirus infection.

5. Conclusion

The study observed that the use of CAM for COVID-19 infection prevention was common with demographic and illness perceptions

towards COVID-19, contributing to the use. The information could be used to educate the safe use of CAM while promoting research into the clinical and potential usefulness for the prevention and treatment of the COVID-19 infection.

6. Availability of data and materials

The data that support the findings of this study are available from the corresponding author, upon reasonable request.

Funding

The project was self-funded by the authors.

CRediT authorship contribution statement

Irene A. Kretchy: Conceptualization, Methodology, Formal analysis, Investigation, Writing – original draft, Writing – review & editing, Project administration. **Joseph A. Boadu:** Methodology, Investigation, Writing – original draft, Writing – review & editing. **James-Paul Kretchy:** Conceptualization, Investigation, Writing – original draft, Writing – review & editing, Project administration. **Kofi Agyabeng:** Methodology, Formal analysis, Writing – original draft, Writing – review & editing. **Alfred A. Passah:** Formal analysis, Writing – review & editing. **Augustina Koduah:** Methodology, Formal analysis, Investigation, Writing – review & editing. **Kwabena F.M. Opuni:** Conceptualization, Methodology, Formal analysis, Investigation, Writing – original draft, Writing – review & editing, Project administration.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Appendix A

Table A1
Plants used as raw materials/homemade preparations for COVID-19 prevention.

	Common name	Scientific name
1	Clove	<i>Syzygium aromaticum</i>
2	Pineapple	<i>Ananas comosus</i>
3	Activated charcoal powder	n.a.
4	Dandelion	<i>Taraxacum officinale</i>
5	Cashew	<i>Anacardium occidentale</i>
6	Pawpaw	<i>Carica papaya</i>
7	Lemon	<i>Citrus Limonum</i>
8	Cinnamon	<i>Cinnamomum zeylanicum</i>
9	Lemongrass	<i>Cymbopogon citratus</i>
10	Ginger	<i>Zingiber officinale</i>
11	Chilli pepper	<i>Capsicum frutescens</i>
12	Neem	<i>Azadirachta indica</i>
13	Garlic	<i>Allium sativum</i>
14	Onion	<i>Allium cepa</i>
15	Negro pepper	<i>Xylopiya aethiopia</i>
16	Grains of paradise	<i>Aframomum melegueta</i>
17	Cloves	<i>Syzygium aromaticum</i>
18	Hibiscus	<i>Hibiscus rosa-sinensis</i>
19	Moringa leaves	<i>Moringa oleifera</i>
20	Prekese	<i>Tetrapleura tetraptera</i>
21	Bitter root	<i>Lewisia rediviva</i>
22	Mango	<i>Mangifera indica</i>
23	African basil	<i>Ocimum gratissimum</i>
24	Pear	<i>Pyrus pyrifolia</i>
25	Turmeric	<i>Curcuma longa</i>
26	Black pepper	<i>Piper guineense</i>

n.a., not applicable

References

- 2010 Population & Housing Census: National Analytical Report https://www.statsghana.gov.gh/gssmain/storage/img/marqueeupdate/Census0_Summary_report_of_final_results.pdf (Accessed on 15th July, 2021).
- Ahmad, Mehmood, Beg, Bilal Mahmood, Majeed, Arfa, Areej, Sadaf, Rifat, Suaaleha, Rasheed, Muhammad Adil, Mahmood, Sammina, Mushtaq, Rana Muhammad Zahid, Hafeez, Mian Abdul, 2021. Epidemiological and clinical characteristics of COVID-19: a retrospective multi-center study in Pakistan. *Front. Public Health* 9. <https://doi.org/10.3389/fpubh.2021.644199>.
- Ahmad, Tauseef, Khan, Muhammad, Haroon, Taha Hussein Musa, Nasir, Saima, Hui, Jin, Bonilla-Aldana, D. Katherine, Rodriguez-Morales, Alfonso J., 2020. COVID-19: Zoonotic aspects. *Travel Med. Infect. Dis.* 36, 101607. <https://doi.org/10.1016/j.tmaid.2020.101607>.
- Aliakbari Dehkordi, M., Eisazadeh, F., Aghajanzadeh, S., 2020. Psychological consequences of patients with coronavirus (COVID-19): A Qualitative Study. *Iran J. Health Psychol.* 2 (2), 9–20. <https://doi.org/10.30473/IJOHP.2020.52395.1074>.
- Alwhaibi, Monira, Bhattacharya, Rituparna, Sambamoorthi, Usha, 2015. Type of multimorbidity and complementary and alternative medicine use among adults. *Evid. Based Complement Alternat. Med.* 2015, 1–9. <https://doi.org/10.1155/2015/362582>.
- Alwhaibi, Monira, Sambamoorthi, Usha, 2016. Sex Differences in the Use of Complementary and Alternative Medicine among Adults with Multiple Chronic Conditions. *Evid. Based Complement Alternat. Med.* 2016, 1–8. <https://doi.org/10.1155/2016/2067095>.
- Alyami, Hamad S., Orabi, Mohamed A.A., Aldhabbah, Fahad M., Alturki, Hamad N., Aburas, Wejdan I., Alfayez, Abdulrahman I., Alharbi, Abdullah S., Almasuod, Rawan A., Alsuhaibani, Njood A., 2020. Knowledge about COVID-19 and beliefs about and use of herbal products during the COVID-19 pandemic: A cross-sectional study in Saudi Arabia. *Saudi Pharm. J.* 28 (11), 1326–1332. <https://doi.org/10.1016/j.jps.2020.08.023>.
- Alyami, M., Serlachius, A., O'Donovan, C.E., van der Werf, B., Broadbent, E., 2021. A systematic review of illness perception interventions in type 2 diabetes: Effects on glycaemic control and illness perceptions. *Diabet. Med.* 38 (3), e14495. <https://doi.org/10.1111/dme.14495>.
- Anakwa, Nella O., Teye-Kwadjo, Enoch, Kretchy, Irene A., 2021. Illness perceptions, social support and antiretroviral medication adherence in people living with HIV in the greater Accra region, Ghana. *Nurs. Open* 8 (5), 2595–2604. <https://doi.org/10.1002/nop2.v8.510.1002/nop2.797>.
- Bishop, F.L., Lewith, G.T., 2010. Who Uses CAM? A Narrative Review of Demographic Characteristics and Health Factors Associated with CAM Use. *Evid. Based Complement Alternat. Med.* 7 (1), 11–28. <https://doi.org/10.1093/ecam/nen023>.
- Boozari, M., Hosseinzadeh, H., 2020. Natural products for COVID-19 prevention and treatment regarding to previous coronavirus infections and novel studies. *Phytother. Res.* 35, 864–876. <https://doi.org/10.1002/ptr.6873>.
- Broadbent, E., Petrie, K.J., Main, J., Weinman, J., 2006. The brief illness perception questionnaire. *J. Psychosom. Res.* 60 (6), 631–637. <https://doi.org/10.1016/j.jpsychores.2005.10.020>.
- Broadbent, E., Wilkes, C., Koschwanez, H., Weinman, J., Norton, S., Petrie, K.J., 2015. A systematic review and meta-analysis of the Brief Illness Perception Questionnaire. *Psychol. Health* 30 (11), 1361–1385. <https://doi.org/10.1080/08870446.2015.1070851>.
- Canaway, R., Manderson, L., 2013. Quality of life, perceptions of health and illness, and complementary therapy use among people with type 2 diabetes and cardiovascular disease. *J. Altern. Complement Med.* 19 (11), 882–890. <https://doi.org/10.1089/acm.2012.0617>.
- Chong, Yuen Yu, Chien, Wai Tong, Cheng, Ho Yu, Chow, Ka Ming, Kassianos, Angelos P., Karekla, Maria, Glover, Andrew, 2020. The role of illness perceptions, coping, and self-efficacy on adherence to precautionary measures for COVID-19. *Int. J. Environ. Res. Public Health* 17 (18), 6540. <https://doi.org/10.3390/ijerph17186540>.
- Chong, Y.Y., Chien, W.T., Cheng, H.Y., Kassianos, A.P., Glover, A.T., Karekla, M., 2021. Can psychological flexibility and prosociality mitigate illness perceptions toward COVID-19 on mental health? A cross-sectional study among Hong Kong adults. *Global Health* 17 (1), 1–13. <https://doi.org/10.1186/s12992-021-00692-6>.
- COVID-19 Dashboard by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University <https://origin-coronavirus.jhu.edu/map.html> (Accessed on September 28, 2021).
- Debas, H.T., Laxminarayan, R., Straus, S.E., 2006. Complementary and alternative medicine. In: *Disease control priorities in developing countries*. Volume 2, ed.: World Bank, Washington, DC, USA; pp. 1281–1292.
- Dias Neto, David, Nunes da Silva, Ana, Roberto, Magda Sofia, Lubenko, Jelena, Constantinou, Marios, Nicolaou, Christina, Lammisios, Demetris, Papacostas, Savvas, Höfer, Stefan, Presti, Giovambattista, Squatrito, Valeria, Vasilioi, Vasilis S., McHugh, Louise, Monestès, Jean-Louis, Baban, Adriana, Alvarez-Galvez, Javier, Paez-Blarrina, Marisa, Montesinos, Francisco, Valdivia-Salas, Sonsoles, Ori, Dorotyya, Lappalainen, Raimo, Kleszcz, Bartosz, Glover, Andrew, Karekla, Maria, Kassianos, Angelos P., 2021. Illness Perceptions of COVID-19 in Europe: Predictors, Impacts and Temporal Evolution. *Front. Psychol.* 12 <https://doi.org/10.3389/fpsyg.2021.6409551>.
- Egede, L.E., Ye, X., Zheng, D., Silverstein, M.D., 2002. The prevalence and pattern of complementary and alternative medicine use in individuals with diabetes. *Diabetes Care* 25 (2), 324–329. <https://doi.org/10.2337/diacare.25.2.324>.
- Erfani, A., Shahriarirad, R., Ranjbar, K., Mirahmadizadeh, A., Moghadami, M., 2020. Knowledge, attitude and practice toward the novel coronavirus (COVID-19) outbreak: a population-based survey in Iran. *Bull. World Health Organ.* 30 (10.2471) <https://doi.org/10.2471/BLT.20.256651>.
- Ernst, E., Fugh-Berman, A., 2002. Complementary and alternative medicine: what is it all about? *Occup. Environ. Med.* 59 (2), 140–144. <https://doi.org/10.1136/oem.59.2.140>.
- Figueiras, M.J., Neto, D.D., 2019. Challenges in “Tailoring” Adjustment. *Eur. Psychol.* 24 (1), 1–6. <https://doi.org/10.1027/1016-9040/a000348>.
- Foxwell, R., Morley, C., Frizelle, D., 2013. Illness perceptions, mood and quality of life: a systematic review of coronary heart disease patients. *J. Psychosom. Res.* 75 (3), 211–222. <https://doi.org/10.1016/j.jpsychores.2013.05.003>.
- Freckleton, Qc I., 2020. COVID-19: Fear, quackery, false representations and the law. *Int. J. Law Psychiatry* 72, 101611. <https://doi.org/10.1016/j.ijlp.2020.101611>.
- Gasmi, A., Chirumbolo, S., Peana, M., Noor, S., Menzel, A., Dadar, M., et al., 2021. The role of diet and supplementation of natural products in COVID-19 prevention. *Biol. Trace Elem. Res.* 1–4 <https://doi.org/10.1007/s12011-021-02623-3>.
- Ghana's Outbreak Response Management Updates <https://ghanahealthservice.org/covid19/archive.php> (Accessed 4th July, 2021).
- Gorbalenya, A.E., Baker, S.C., Baric, R.S., de Groot, R.J., Drosten, C., Gulyaeva, A.A., et al., 2020. The species Severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. *Nat. Microbiol.* 5 (4), 536–544. <https://doi.org/10.1038/s41564-020-0695-z>.
- Gyasi, Razak Mohammed, Asante, Felix, Abass, Kabila, Yeboah, Joseph Yaw, Adu-Gyamfi, Samuel, Amoah, Padmore Adusei, Halsall, Jamie, 2016. Do health beliefs explain traditional medical therapies utilisation? Evidence from Ghana. *Cogent. Sci. Soc.* 2 (1), 1209995. <https://doi.org/10.1080/23311886.2016.1209995>.
- Gyasi, R.M., Siaw, L.P., Mensah, C.M., 2015. Prevalence and pattern of traditional medical therapy utilisation in Kumasi metropolis and Sekyere south district, Ghana. *J. Ethnopharmacol.* 161, 138–146. <https://doi.org/10.1016/j.jep.2014.12.004>.
- Gyasi, R.M., Mensah, C.M., Siaw, L.P., 2015. Predictors of traditional medicines utilisation in the Ghanaian health care practice: interrogating the Ashanti situation. *J. Community Health* 40 (2), 314–325. <https://doi.org/10.1007/s10900-014-9937-4>.
- Haines, C., Loades, M., Davis, C., 2019. Illness perceptions in adolescents with chronic fatigue syndrome and other physical health conditions: Application of the common sense model. *Clin. Child. Psychol. Psychiatry* 24 (3), 546–563. <https://doi.org/10.1111/1359104519829796>.
- Harris, P.E., Cooper, K.L., Relton, C., Thomas, K.J., 2012. Prevalence of complementary and alternative medicine (CAM) use by the general population: a systematic review and update. *Int. J. Clin. Pract.* 66 (10), 924–939. <https://doi.org/10.1111/j.1742-1241.2012.02945.x>.
- Hossain, Mohammad Bellal, Alam, Md. Zakiul, Islam, Md. Syful, Sultan, Shafayat, Faysal, Md. Mahir, Rima, Sharmin, Hossain, Md. Anwer, Mahmood, Maliha Mubashirah, Kashfi, Shaima Shohuda, Mamun, Abdullah al, Monia, Hossna Tasmia, Shoma, Sharmin Sultana, 2020. Do knowledge and attitudes matter for preventive behavioral practices toward the COVID-19? A cross-sectional online survey among the adult population in Bangladesh. *Heliyon* 6 (12), e05799. <https://doi.org/10.1016/j.heliyon.2020.e05799>.
- Jabaris, Sugin Lal S., Ananthakshmi, V., 2021. The current situation of COVID-19 in India. *Brain Behav. Immun. Health* 11, 100200. <https://doi.org/10.1016/j.bbih.2021.100200>.
- James, Peter Bai, Wardle, Jon, Steel, Amie, Adams, Jon, 2018. Traditional, complementary and alternative medicine use in Sub-Saharan Africa: a systematic review. *BMJ Glob. Health* 3 (5), e000895. <https://doi.org/10.1136/bmjgh-2018-00089510.1136/bmjgh-2018-000895.suppl110.1136/bmjgh-2018-000895.suppl2>.
- Kretchy, I.A., Asiedu-Danso, M., Kretchy, J.-P., 2021. Medication management and adherence during the COVID-19 pandemic: perspectives and experiences from low- and middle-income countries. *Res. Social Adm. Pharm.* 17 (1), 2023–2026. <https://doi.org/10.1016/j.sapharm.2020.04.007>.
- Kretchy, Irene A., Koduah, Augustina, Opuni, Kwabena F.M., Agyabeng, Kofi, Ohene-Agyei, Thelma, Bofo, Eugene A., Ntwi, Prisca O., 2021. Prevalence, patterns and beliefs about the use of herbal medicinal products in Ghana: a multi-centre community-based cross-sectional study. *Trop. Med. Int. Health* 26 (4), 410–420. <https://doi.org/10.1111/tmi.v26.410.1111/tmi.13541>.
- Kretchy, I.A., Owusu-Daaku, F., Danquah, S., 2014. Patterns and determinants of the use of complementary and alternative medicine: a cross-sectional study of hypertensive patients in Ghana. *BMC Complement Altern. Med.* 14 (1), 1–7. <https://doi.org/10.1186/1472-6882-14-44>.
- Kristoffersen A, van der Werf E, Stub T, Musial F, Wider B, Jong M, Wode K, Danell JA, Busch M, Hoenders HR, Nordberg J. Consultations with Health Care Providers and Use of Self-management Strategies for Prevention and Treatment of COVID-19 Related Symptoms. A Population Based Cross-sectional Study in Norway, Sweden and the Netherlands. <https://www.researchsquare.com/article/rs-145141/v1>.
- Kristoffersen, A.E., Stub, T., Salamonsen, A., Musial, F., Hamberg, K., 2014. Gender differences in prevalence and associations for use of CAM in a large population study. *BMC Complement Altern. Med.* 14, 463. <https://doi.org/10.1186/1472-6882-14-463>.
- Kucuk, E.E., 2017. Assessment of the use of complementary and alternative medicine with respect to illness perception among individuals with chronic diseases. *Med. Sci.* 6 (3), 457–463. <https://doi.org/10.5455/medscience.2017.06.8592>.
- Kuunien, N., Domanban, P.B., 2012. Demand for complementary and alternative medicine in Ghana. *Int. J. Humanit. Soc. Sci.* 2, 288–294.
- Laiyemo, M.A., Nunlee-Bland, G., Adams, R.G., Laiyemo, A.O., Lombardo, F.A., 2015. Characteristics and health perceptions of complementary and alternative medicine users in the United States. *Am. J. Med. Sci.* 349 (2), 140–144. <https://doi.org/10.1097/MAJ.0000000000000363>.
- Leventhal, H., Meyer, D., Nerenz, D., 1980. The Common-Sense Representation of Illness Danger. In: *Contributions to Medical Psychology*. Edited by Rachman S, vol. 2: New York: Pergamon Press; pp. 7–30.

- Leventhal, H., Phillips, L.A., Burns, E., 2016. The Common-Sense Model of Self-Regulation (CSM): a dynamic framework for understanding illness self-management. *J. Behav. Med.* 39 (6), 935–946. <https://doi.org/10.1007/s10865-016-9782-2>.
- Lichtenstein, G., Waalen, J., 2002. Distinguishing complementary medicine from alternative medicine. *Arch. Intern. Med.* 162 (8), 943.
- Luo, Hui, Tang, Qiao-ling, Shang, Ya-xi, Liang, Shi-bing, Yang, Ming, Robinson, Nicola, Liu, Jian-ping, 2020. Can Chinese medicine be used for prevention of corona virus disease 2019 (COVID-19)? A review of historical classics, research evidence and current prevention programs. *Chin. J. Integr. Med.* 26 (4), 243–250. <https://doi.org/10.1007/s11655-020-3192-6>.
- Man, Milena Adina, Toma, Claudia, Motoc, Nicoleta Stefania, Necrelescu, Octavia Luiza, Bondor, Cosmina Ioana, Chis, Ana Florica, Lesan, Andrei, Pop, Carmen Monica, Todea, Doina Adina, Dantes, Elena, Puiu, Ruxandra, Rajnoveanu, Ruxandra-Mioara, 2020. Disease perception and coping with emotional distress during COVID-19 pandemic: a survey among medical staff. *Int. J. Environ. Res. Public Health* 17 (13), 4899. <https://doi.org/10.3390/ijerph17134899>.
- Mentis, Alexios-Fotios A., 2021. What neurological and psychiatric effects does COVID-19 have on its survivors? *BMC Med.* 19 (1) <https://doi.org/10.1186/s12916-021-01986-7>.
- National Health Commission of the People's Republic of China. Diagnosis and treatment protocol for COVID-19 (trial version 7). http://en.nhc.gov.cn/2020-03/29/c_78469.htm. Accessed September 28, 2021.
- Nekliudov, Nikita A, Blyuss, Oleg, Cheung, Ka Yan, Petrou, Loukia, Genuneit, Jon, Sushentsev, Nikita, Levadnaya, Anna, Comberiat, Pasquale, Warner, John O, Tudor-Williams, Gareth, Teufel, Martin, Greenhawt, Matthew, DunnGalvin, Audrey, Munblit, Daniel, 2020. Excessive media consumption about COVID-19 is associated with increased state anxiety: outcomes of a large online survey in Russia. *J. Med. Internet Res.* 22 (9), e20955. <https://doi.org/10.2196/20955>.
- Okai, G.A., Abekah-Nkrumah, G., Asuming, P.O., 2019. Perceptions and trends in the use of community pharmacies in Ghana. *J. Pharm. Policy Pract.* 12 (1), 1–9. <https://doi.org/10.1186/s40545-019-0186-x>.
- Okoronkwo, Ijeoma, Onyia-pat, Jane-lovena, Okpala, Pat, Agbo, Mary-Ann, Ndu, Afam, 2014. Patterns of complementary and alternative medicine use, perceived benefits, and adverse effects among adult users in Enugu Urban, Southeast Nigeria. *Evid. Based Complement Alternat. Med.* 2014, 1–6. <https://doi.org/10.1155/2014/239372>.
- Panyod, S., Ho, C.-T., Sheen, L.-Y., 2020. Dietary therapy and herbal medicine for COVID-19 prevention: A review and perspective. *J. Tradit. Complement Med.* 10 (4), 420–427. <https://doi.org/10.1016/j.jtcm.2020.05.004>.
- Paudyal, V., Sun, S., Hussain, R., Abutaleb, M.H., Hedima, E.W., 2021. Complementary and alternative medicines use in COVID-19: A global perspective on practice, policy and research. *Res. Social Adm. Pharm.* <https://doi.org/10.1016/j.sapharm.2021.05.004>.
- Piltch-Loeb, Rachael, Merdjanoff, Alexis, Meltzer, Gabriella, 2021. Anticipated mental health consequences of COVID-19 in a nationally-representative sample: Context, coverage, and economic consequences. *Prev. Med.* 145, 106441. <https://doi.org/10.1016/j.ypmed.2021.106441>.
- National Clinical Management Protocol based on Ayurveda and Yoga for management of Covid-19. <https://www.ayush.gov.in/docs/ayush-Protocol-covid-19.pdf>. Accessed September 28, 2021.
- Pullen, M.F., Skipper, C.P., Hullsiek, K.H., Bangdiwala, A.S., Pastick, K.A., Okafor, E.C., et al., 2020. Symptoms of COVID-19 Outpatients in the United States. *Open Forum. Infect. Dis.* 7 (7) <https://doi.org/10.1093/ofid/ofaa271>.
- Sanders, J.M., Monogue, M.L., Jodlowski, T.Z., Cutrell, J.B., 2020. Pharmacologic treatments for coronavirus disease 2019 (COVID-19): a review. *JAMA* 323 (18), 1824–1836. <https://doi.org/10.1001/jama.2020.6019>.
- Silveira, Damaris, Prieto-Garcia, Jose Maria, Boylan, Fabio, Estrada, Omar, Fonseca-Bazzo, Yris Maria, Jamal, Claudia Masrouah, Magalhães, Pérola Oliveira, Pereira, Edson Oliveira, Tomczyk, Michal, Heinrich, Michael, 2020. COVID-19: is there evidence for the use of herbal medicines as adjuvant symptomatic therapy? *Front. Pharmacol.* 11 <https://doi.org/10.3389/fphar.2020.581840>.
- Singh, V., Raidoo, D.M., Harries, C.S., 2004. The prevalence, patterns of usage and people's attitude towards complementary and alternative medicine (CAM) among the Indian community in Chatsworth, South Africa. *BMC Complement Altern. Med.* 4 (1), 1–7. <https://doi.org/10.1186/1472-6882-4-3>.
- Skapinakis, Petros, Bellos, Stefanos, Oikonomou, Achilleas, Dimitriadis, Georgios, Gkikas, Paschalis, Perdikari, Evridiki, Mavreas, Venetsanos, 2020. Depression and its relationship with coping strategies and illness perceptions during the COVID-19 lockdown in Greece: a cross-sectional survey of the population. *Depress Res. Treat.* 2020, 1–11. <https://doi.org/10.1155/2020/3158954>.
- Tarighi, Parastoo, Eftekhari, Samane, Chizari, Milad, Sabernavaei, Mahsa, Jafari, Davod, Mirzabeigi, Parastoo, 2021. A review of potential suggested drugs for coronavirus disease (COVID-19) treatment. *Eur. J. Pharmacol.* 895, 173890. <https://doi.org/10.1016/j.ejphar.2021.173890>.
- Valizadeh, R., Dadashzadeh, N., Zakeri, R., Kellner, S.J., Rahimi, M.M., 2020. Drug therapy in hospitalized patients with very severe symptoms following COVID-19. *Med. Sci.* 35, e79. <https://doi.org/10.34172/npj.2020.21>.
- VanderWeele, T.J., 2020. Challenges estimating total lives lost in COVID-19 decisions: consideration of mortality related to unemployment, social isolation, and depression. *JAMA* 324 (5), 445–446. <https://doi.org/10.1001/jama.2020.12187>.
- Wu, Z., McGoogan, J., 2020. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72314 cases from the Chinese Center for Disease Control and Prevention. *JAMA*. <https://doi.org/10.1001/jama.2020.2648>.
- Yang, Rong, Mei, Hui, Zheng, Tongzhang, Fu, Qiang, Zhang, Yiming, Buka, Stephen, Yao, Xinan, Tang, Zezhong, Zhang, Xichi, Qiu, Lin, Zhang, Yaqi, Zhou, Jieqiong, Cao, Jiangxia, Wang, Youjie, Zhou, Aifen, 2020. Pregnant women with COVID-19 and risk of adverse birth outcomes and maternal-fetal vertical transmission: a population-based cohort study in Wuhan, China. *BMC Med.* 18 (1) <https://doi.org/10.1186/s12916-020-01798-1>.
- Yarney, J., Donkor, A., Opoku, S.Y., Yarney, L., Agyeman-Duah, I., Abakah, A.C., et al., 2013. Characteristics of users and implications for the use of complementary and alternative medicine in Ghanaian cancer patients undergoing radiotherapy and chemotherapy: a cross-sectional study. *BMC Complement Altern. Med.* 13 (1), 1–9. <https://doi.org/10.1186/1472-6882-13-16>.
- Yawson, Alfred E., Oduro-Mensah, Ebenezer, Tetteh, John, Adomako, Isaac, Adjei-Mensah, Evelyn, Owoo, Christian, Yawson, Anita O., Oliver-Commey, Joseph A., Pupilamp, Peter, Samba, Ali, Lartey, Margaret, 2020. Clinical features of COVID-19 in Ghana: symptomatology, illness severity and comorbid non-communicable diseases. *Ghana Med. J.* 54 (4s), 23–32. <https://doi.org/10.4314/gmj.v54i4s.5>.
- Zhang, Yan, Leach, Matthew J., Hall, Helen, Sundberg, Tobias, Ward, Lesley, Sibbritt, David, Adams, Jon, 2015. Differences between Male and Female Consumers of Complementary and Alternative Medicine in a National US Population: A Secondary Analysis of 2012 NHIS Data. *Evid. Based Complement Altern. Med.* 2015, 1–10. <https://doi.org/10.1155/2015/413173>.